




Blackburn Bay Condition Report for 2014




PASS



Chl-*a*



N



P

3 out of 3
indicators were
rated as **PASS**.

All three
indicators must pass for the bay to be rated as **PASS**.

Summary:

The overall health of Blackburn Bay has remained good in 2014 with two out of three water quality parameters, chlorophyll *a* and phosphorus, below their associated target levels. Nitrogen concentration remained fairly steady and is in the "Good" range, between target and threshold levels.

Water Quality: All three water quality indicators (chlorophyll *a*, nitrogen, and phosphorus) were rated as pass (below threshold). The mean for chlorophyll *a* was calculated as an arithmetic mean and the means for nitrogen and phosphorus were calculated as geometric means (per the Numeric Nutrient Criteria outlined in the Florida Administrative Code, section 62-302.532). The mean chlorophyll *a* level (0.0043 mg/l) again decreased, keeping it well below the target level of 0.006 mg/l. The mean nitrogen level increased almost imperceptibly from 2013, to 0.4226 mg/l, retaining its "Good" score. Phosphorus concentrations in the bay have remained relatively



Bays included in this report:
Blackburn Bay

constant and low, with a mean of 0.0647 mg/l, well below the target (0.170 mg/l) and threshold (0.210 mg/l) levels and earning a score of "Excellent".

Biotic Indicator: Measurement of the biotic indicator, seagrass, was performed in 2014 by the Southwest Florida Water Management District. Total seagrass acreage in Blackburn Bay was measured to be 344 acres. Seagrass acreage continues to slowly increase, but it is still below the target level of 447 acres.

Water Chemistry Ratings

Total nitrogen, total phosphorus, and chlorophyll *a* levels are monitored carefully by water resource managers and used by regulatory authorities to determine whether a bay meets the water quality standards mandated by the Clean Water Act. The trend graphs for these indicators are shown below, along with their target and threshold values. A target value is a desirable goal to be attained, while a threshold is an undesirable level which is to be avoided. An individual indicator receives an "Excellent" rating if its mean value is below the target, a "Good" rating if its mean value is above the target but does not exceed the threshold, and a "Caution" rating if the mean value exceeds the threshold.


The charts below illustrate the general trend of water quality parameters. They show a six-month running average, which moderates high and low values in the data.



Chlorophyll a

Score: Excellent

Units: ug/l	Year 2014	Historical period of record
High	23.87	43.00
Mean	4.31	4.98
Low	1.09	0.20
No. of Samples	58	902




N

Nitrogen, Total

Score: Good

Units: mg/l	Year 2014	Historical period of record
High	0.855	2.205
Mean	0.423	
Low	0.265	0.055
No. of Samples	58	896




P

Phosphorus, Total

Score: Excellent


Units: mg/l	Year 2014	Historical period of record
High	0.120	0.530
Mean	0.065	0.102
Low	0.050	0.050
No. of Samples	58	912

**Other Measures of Bay Health**

In addition to nutrient levels and chlorophyll concentration, dissolved oxygen levels, and water clarity are also objective indicators of bay health. These have complex interactive cycles which are affected by rainfall, temperature, and tidal action, as well as other factors. High nutrient levels (nitrogen and phosphorus) can stimulate excessive growth of marine algae (indicated by chlorophyll *a* level), resulting in reduced water clarity (and increased light attenuation) and depleted oxygen levels. Both plants and animals in a bay need oxygen to survive, and the seagrasses which provide food and cover for bay creatures need light for photosynthesis.


Dissolved Oxygen

Units: mg/l	Year 2014	Historical period of record
High	8.20	11.90
Mean	6.00	6.47
Low	3.20	1.60
No. of Samples	48	956



Apparent Color

Units: PCU	Year 2014	Historical period of record
High	70.00	250.00
Mean	11.79	18.97
Low	2.00	2.00
No. of Samples	58	912




BOD, Biochemical oxygen demand

Units: mg/l	Year 2014	Historical period of record
High	1.90	7.10
Mean	0.70	1.07
Low	0.50	0.50
No. of Samples	58	801




Dissolved oxygen saturation

Units: percent (%)	Year 2014	Historical period of record
High	115.00	198.00
Mean	87.29	94.69
Low	51.00	26.00
No. of Samples	48	956




Karenia brevis ("red tide")

Units: #/l	Year 2014	Historical period of record
High	1000.00	2280000.00
Mean	1000.00	35570.08
Low	1000.00	1000.00
No. of Samples	58	528



Light Attenuation

Units: K(1/m)	Year 2014	Historical period of record
High	2.31	5.03
Mean	0.75	0.86
Low	0.35	0.16
No. of Samples	47	794




Nitrogen, Ammonia + Ammonium as N

Units: ug/l	Year 2014	Historical period of record
High	67.00	120.00
Mean	16.02	15.35
Low	5.00	5.00
No. of Samples	58	914




Nitrogen, Kjeldahl

Units: ug/l	Year 2014	Historical period of record
High	850.00	2200.00
Mean	429.83	346.07
Low	260.00	0.05
No. of Samples	58	913




Nitrogen, Nitrite + Nitrate as N

Units: ug/l	Year 2014	Historical period of record
High	8.00	65.00
Mean	5.41	6.00
Low	5.00	5.00
No. of Samples	58	1234




pH

Units: None	Year 2014	Historical period of record
High	8.20	8.40
Mean	7.99	7.94
Low	7.80	6.10
No. of Samples	48	956




Salinity

Units: PSS	Year 2014	Historical period of record
High	36.20	39.30
Mean	33.07	33.25
Low	22.80	5.70
No. of Samples	48	956




Specific conductance

Units: umho	Year 2014	Historical period of record
High	54520.00	58760.00
Mean	50341.46	50563.91
Low	36150.00	10130.00
No. of Samples	48	956




Temperature, water

Units: deg F	Year 2014	Historical period of record
High	86.90	92.84
Mean	76.53	76.41
Low	65.84	47.84
No. of Samples	48	956



Turbidity

Units: NTU	Year 2014	Historical period of record
High	30.00	39.00
Mean	3.81	3.96
Low	1.20	0.20
No. of Samples	58	903




Annual Averages

Indicator	Units	2010	2011	2012	2013	2014	Trend
Dissolved Oxygen	mg/l	6.92	6.41	6.11	6.11	6.00	
Dissolved oxygen saturation	percent (%)	95.81	92.48	89.71	87.79	87.29	
Light Attenuation	K(1/m)	0.92	0.75	0.88	0.91	0.75	
Salinity	PSS	32.60	33.44	33.19	31.31	33.07	
Turbidity	NTU	5.08	4.20	3.83	4.06	3.81	

Bay Contour Maps (2014)

Contour mapping is one of the best ways to visualize spatial differences in coastal water quality. The interactive map shown below presents monthly data for one selected water quality indicator atop an aerial view of the bay. Choose a different water quality parameter from the list at the top to change the map.

Showing 2014 Monthly Contour Maps for: Chlorophyll a
January




Contour Legend:

- Less than 1 mg/l
- 1.0 - 5.9 mg/l
- 6.0 - 10.9 mg/l
- 11.0 - 17.9 mg/l
- Greater than 18 mg/l


Seagrasses

Among the most important habitats in Florida's estuarine environments, seagrass beds are indispensable for the role they play in cycling nutrients, supplying food for wildlife, stabilizing sediments, and providing habitat for juvenile and adult finfish and shellfish. Use the interactive map below to observe the size, density and location of seagrass beds from year to year. The graph shows how the total amount of seagrass in the bay has changed over time. Seagrass calculations are aggregates of patchy and continuous seagrass measurements only. Recordings of attached algae are not included in these summaries.

Showing Seagrass Coverage for 2020:



Seagrass Acreage Variation within Blackburn Bay



Impervious Features

Rain that falls on land that is in a natural state is absorbed and filtered by soils and vegetation as it makes its way into underground aquifers. However, in developed areas, "impervious surfaces" impede this process and contribute to polluted urban runoff entering surface waters. These surfaces include human infrastructure like roads, sidewalks, driveways and parking lots that are covered by impenetrable materials such as asphalt, concrete, brick and stone, as well as buildings and other permanent structures. Soils that have been disturbed and compacted by urban development are often impervious as well.




14% of the land area within the **Little Sarasota Bay Watershed** is covered by

impervious surfaces

2014 Impervious Surface Coverage by Type

in acres, within the Little Sarasota Bay Watershed











Land Use / Land Cover

Land use within a bay's watershed has a major effect on its water quality. In general, less development means better water quality. Land Cover/Land Use classifications categorize land in terms of its observed physical surface characteristics (upland or wetland, e.g.), and also reflect the types of activity that are taking place on it (agriculture, urban/built-up, utilities, etc.). Florida uses as its standard a set of statewide classifications which were developed by the Florida Department of Transportation.

Blackburn Bay is located within the Little Sarasota Bay Watershed. The chart below shows the land use / land cover characteristics for Little Sarasota Bay Watershed within the boundary of this Water Atlas. **[View details about the Little Sarasota Bay Watershed »](#)**

Acreege and Percentage within each Land Use / Land Cover Category for Little Sarasota Bay Watershed

2014 Bay Conditions Report for Blackburn Bay

Land Use Classification	1990	2005	2011	2014	2017	2020	Trend
Urban & Built-up	8,943 31.9%	11,834 42.2%	12,102 43.1%	12,162 43.3%	12,777 45.5%	13,343 47.5%	
Agriculture	3,550 12.6%	3,228 11.5%	3,258 11.6%	4,223 15%	3,124 11.1%	2,837 10.1%	
Rangeland	825 2.9%	1,822 6.5%	1,474 5.3%	579 2.1%	1,233 4.4%	877 3.1%	
Upland Forests	7,098 25.3%	3,066 10.9%	2,981 10.6%	2,725 9.7%	2,687 9.6%	2,603 9.3%	
Water	3,429 12.2%	4,123 14.7%	4,147 14.8%	4,175 14.9%	4,227 15.1%	4,337 15.5%	
Wetlands	3,490 12.4%	3,133 11.2%	3,191 11.4%	3,227 11.5%	3,121 11.1%	3,120 11.1%	
Barren Land	62 0.2%	18 0.1%	19 0.1%	20 0.1%	6 0%	6 0%	
Transportation and Utilities	675 2.4%	841 3%	892 3.2%	952 3.4%	898 3.2%	948 3.4%	

2020 Land Use / Land Cover for Little Sarasota Bay Watershed

as a percentage of land area for this watershed

